

**Application
for
United States Letters Patent**

RECEIVED
MAR 12 1993

To all whom it may concern:

Be it known that Cy A. Stein

have invented certain new and useful improvements in
OLIGONUCLEOTIDE INHIBITORS OF bcl-xL

of which the following is a full, clear and exact description.

OLIGONUCLEOTIDE INHIBITORS OF bcl-xL

This application is a continuation-in-part of U.S. Serial No. 09/109,614, filed July 2, 1998, the content of which is hereby incorporated by reference.

Throughout this application, various references are referred to. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.

Background of the Invention

Bcl-xL is an important anti-apoptotic protein that belongs to the bcl-2 family. Bcl-xL is a critical determinant of intimal lesion formation and thus is an important contributor to the progression of vascular disease. Pollman et al. (1998) Nature Medicine 4: 222-227. In addition, bcl-xL has been implicated as a causative factor in cancer.

The subject invention involves oligonucleotides that reduce or eliminate the expression of bcl-xL. Synthetic oligodeoxynucleotides have been utilized as antisense inhibitors of mRNA translation in vitro and in vivo. Beaucage, S., and Caruthers, M., (1981) Tetrahedron Lett. 37:3557; Iverson, P. (1991) Anti-Cancer Drug Des. 6:531; Ratajczak, et al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89:11823; Uhlmann, E., and Peyman, A. (1990) Chem. Rev. 90:544-579. Antisense oligonucleotides have found widespread application because of their abilities to control and/or inhibit gene expression in a selective manner in cellular systems. Ghosh, S., et al. (1990) J. Biol. Chem. 265:2935-2940; Hemken, P., et al. (1992) J. Biol. Chem. 267:9948-0057; Lestinger, R., U.S. Patent No. 4,958,103, issued September 18, 1990; Shewmaker et al.,

U.S. Patent no. 5,107,065, issued April 21, 1992; Tullis,
U.S. Patent No., issued June 11, 1991; Zhao, Q., et al.
(1993) Antisense Research and Development 3:53-66. Thus,
5 the subject invention represents a precise and selective
way of regulating bcl-xL expression both in vivo and in
vitro.

Summary of the Invention

This invention provides an antisense oligonucleotide or analog thereof comprising 10 or more contiguous bases or base analogs from the sequence of bases of sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

This invention also provides the above-described antisense oligonucleotides, wherein the nucleotide sequence comprises nucleotide sequence A, A', B, C, C', D, E, E', F, G, G', H, H', I, I', J, K, K', L, L', M, or M' of Figures 2A and 2B.

This invention also provides the above-described antisense oligonucleotides, wherein the oligonucleotide is encapsulated in a liposome or nanoparticle.

This invention also provides the above-described antisense oligonucleotides, wherein the phosphate backbone comprises phosphorothioate bonds.

In addition, this invention provides a method of treating cancer, comprising introducing into a tumor cell an effective amount of the the above-described antisense oligonucleotide, thereby reducing the levels of bcl-xL protein produced and treating cancer.

This invention further provides a method of treating cancer, comprising introducing into a tumor cell an effective amount of the the above-described antisense oligonucleotide, thereby reducing the levels of bcl-2 protein produced and treating cancer.

This invention also provides the above-described methods, wherein the introducing comprises using amine and cationic delivery reagents.

This invention also provides the above-described methods, wherein the introducing comprises using porphyrin or lipofectin as a delivery agent.

This invention further provides the above-described pharmaceutical compositions, wherein the pharmaceutical composition comprises tetra meso-(4-methylpyridyl)porphine or tetra meso-(anilinium)porphine or a combination thereof.

Brief Description of the Figures

Figure 1

Oligonucleotide sequences complementary to bcl-X_L mRNA.

Figures 2A and 2B

5 Oligonucleotide sequences and analogs thereof complementary to bcl-X_L mRNA.

Figures 3A, 3B, 3C, and 3D

Some of the many possible phosphodiester moiety analogs.

Figure 4

10 An example of a sugar containing an aminoalkyloxy linker.

Figures 5A and 5B

Some of the many possible base moiety analogs.

Figure 6

15 Effect of 18-mer PS oligonucleotides on bcl-xL protein expression in LNCaP cells. The height of columns represents oligonucleotide inhibition of bcl-xL protein expression. Oligonucleotides were added to cells at a concentration of 1 μ M in the presence of 3 μ M of TAP for 4 h. The most active compound was T31062.

20 Figure 7

Western blot analysis of bcl-xL protein expression in prostate cancer cell lines after oligonucleotide treatment. The treatment of the cell with complex of oligonucleotides T31057, T31061, T31062 (1 μ M) with TMP (2 μ M) or TAP (3 μ M) down-regulates bcl-xL protein expression, whereas oligonucleotides alone do not cause this effect.

25

Figure 8

Antisense Oligo nucleotides to Bcl-xL mRNA

Figure 9

30 Regulation of Bcl-Family Proteins with 2"-O-Methyl -

Modified PS Oligonucleotides in T24 Cell Line

Figure 10

Most active chimeric PS-PO oligonucleotides by their ability to down-regulate Bcl-xL protein expression

5 Figure 11

Regulation of Bcl-family proteins with 2'-O-Methyl-Modified PS Oligonucleotides in PC - 3 Cell Line

Figure 12

10 Regulation of Bcl-xL and Bax proteins with 2'-O-Methyl-Modified PS oligonucleotides in LNCaP Cell Line

Figure 13

Down-regulation of Bcl-family proteins expression with PS-PO oligonucleotides in LNCaP cell line

Figure 14

15 Down-regulation of Bcl-family proteins expression with PS-PO oligonucleotides in PC3 cell line

Figure 15

Regulation of Bcl-Family Proteins with PS-PO oligonucleotides in T24 Cell line

20 Figure 16

Down-regulation of Bcl-xL mRNA with aPS-PO oligonucleotides in T24 cell line

Figure 17

25 Regulation of Bcl-Family Proteins with PS-PO oligonucleotides in LNCaP cell line

Detailed Description of the Invention

The following standard abbreviations are used throughout the specification to indicate specific nucleotides:

5 C = Cytidine A = Adenosine
 T = Thymidine G = Guanosine

10 Nucleic acid synthesizers are available to synthesize oligonucleotides of any desired sequence. Certain oligonucleotide analogs may also be readily synthesized by modifying the reactants and reaction conditions. For example, phosphorothioate and methylphosphonate oligonucleotides may be synthesized using commercially available automated oligonucleotide synthesizers.

15 An oligonucleotide's binding affinity to a complementary nucleic acid may be assessed by determining the melting temperature (T_m) of a hybridization complex. The T_m is a measure of the temperature required to separate the nucleic acid strands of a hybridization complex. The T_m may be measured by using the hybridization complex's UV spectrum to assess the degree and strength of
20 hybridization. During hybridization, base stacking occurs which reduces the UV absorption of the nucleic acid. By monitoring UV absorption and the resulting increase in UV absorption that occurs during strand separation, one may assess the hybridization affinity of a nucleic acid for its
25 complement.

The structure and stability of hybridization complexes may be further assessed using NMR techniques known to those skilled in the art.

30 A vast array of oligonucleotide analogs exist that achieve the same functionality as naturally occurring oligonucleotides. There is an extensive literature setting forth an almost limitless variety of modifications that can be used to generate oligonucleotide analogs. The

phosphophate, sugar, and/or base moieties may be modified and/or replaced by the introduction/removal of chemical groups and/or bonds. Many oligonucleotide analogs have superior properties to those of naturally occurring oligonucleotides. Such superior properties include, but are not limited to, increased hybridization affinity and/or resistance to degradation.

Phosphodiester Moiety Analogs

Numerous analogs to the naturally occurring phosphodiester backbone have been used in oligonucleotide design. Phosphorothioate, phosphorodithioate, and methylphosphonate are readily synthesized using known chemical methods. Because novel nucleotide linkages can be synthesized manually to form a dimer and the dimer later introduced into the oligonucleotide via automated synthesis, the range of potential backbone modifications is as broad as the scope of synthetic chemistry. For example, the oligonucleotide may be substituted or modified in its internucleotide phosphate residue with a thioether, carbamate, carbonate, acetamdate or carboxymethyl ester. While all of the backbone modifications that have been characterized are too broad to set forth, Figures 3A, 3B, 3C, and 3D illustrate some of the many backbone modifications that may be used in oligonucleotide analogs.

Unlike the naturally occurring phosphodiester moieties, many phosphodiester analogs have chiral centers. For example, phosphorothioates, methylphosphonates, phosphoramidates, and alkyl phosphotriesters all have chiral centers. One skilled in the art would recognize numerous other phosphodiester analogs that possess chiral centers. Because of the importance of stereochemistry in hybridization, the stereochemistry of phosphodiester analogs can influence the hybridization affinity of the oligonucleotide for its target.

Most phosphodiester backbone analogs exhibit increased

resistance to nuclease degradation. In an embodiment, phosphorothioates, methyl phosphonates, phosphorimidates, and/or phosphotriesters are used to achieve enhanced nuclease resistance. Increased resistance to degradation may also be achieved by capping the 5' and/or 3' end of the oligonucleotide. In an embodiment, the 5' and/or 3' end capping of the oligonucleotide is via a 5'-5' and/or 3'-3' terminal inverted linkage.

Phosphorothioate oligodeoxynucleotides are relatively nuclease resistant, water soluble analogs of phosphodiester oligodeoxynucleotides. These molecules are racemic, but still hybridize well to their RNA targets. Stein, C., et al. (1991) Pharmac. Ther. 52:365-384.

Phosphorothioate oligonucleotides may be stereo regular, stereo non-regular or stereo random. A stereo regular phosphorothioate oligonucleotide is a phosphorothioate oligonucleotide in which all of the phosphodiester linkages or phosphorothiodiester linkages polarize light in the same direction. Each phosphorous in each linkage may be either an Sp or Rp diastereomer.

Sugar Moiety Analogs

Oligonucleotide analogs may be created by modifying and/or replacing a sugar moiety.

The sugar moiety of the oligonucleotide may be modified by the addition of one or more substituents. For example, one or more of the sugar moieties may contain one or more of the following substituents: amino-alkylamino, araalkyl, heteroalkyl, heterocycloalkyl, aminoalkylamino, O, H, an alkyl, polyalkylamino, substituted silyl, F, Cl, Br, CN, CF₃, OCF₃, OCN, O-alkyl, S-alkyl, SO Me, QNO, NH-alkyl, OCH₂CH=CH₂, OCH₂CCH, OCCHO, allyl, O-allyl, NQ₂, N₃, and NH₂.

Modification of the 2' position of the ribose sugar has

been shown in many instances to increase the oligonucleotide's resistance to degradation.

For example, the 2' position of the sugar may be modified to contain one of the following groups: H, OH, OCN, O-alkyl, F, CN, CF₃, allyl, O-allyl, OCF₃, S-alkyl, SO₂Me, ONO₂, NO₂, N₃, NH₂, NH-alkyl, or OCH=CH₂, OCCH, wherein the alkyl may be straight, branched, saturated, or unsaturated.

In addition, the oligonucleotide may have one or more of its sugars modified and/or replaced so as to be a ribose or hexose (i.e. glucose, galactose).

Further, the oligonucleotide may have one or more α -anomeric sugars. The oligonucleotide may also have one or more L sugars.

The sugar may be modified to contain one or more linkers for attachment to other chemicals such as fluorescent labels. In an embodiment, the sugar is linked to one or more aminoalkyloxy linkers. An example of a sugar containing an aminoalkyloxy linker is shown in Figure 4. In another embodiment, the sugar contains one or more alkylamino linkers. Aminoalkyloxy and alkylamino linkers may be attached to biotin, cholic acid, fluorescein, or other chemical moieties through their amino group.

Base Moiety Analogs

In addition, the oligonucleotide may have one or more of its nucleotide bases substituted or modified. In addition to adenine, guanine, cytosine, thymine, and uracil, other bases such as inosine, deoxyinosine, hypoxanthine may be used. In addition, isoteric purine 2'-deoxy-furanoside analogs, 2'-deoxynebularine or 2'-deoxyxanthosine, or other purine or pyrimidine analogs may also be used. By carefully selecting the bases and base analogs, one may fine tune the hybridization properties of the

oligonucleotide. For example, inosine may be used to reduce hybridization specificity, while diaminopurines may be used to increase hybridization specificity.

5 Adenine and guanine may be modified at positions N3, N7, N9, C2, C4, C5, C6, or C8 and still maintain their hydrogen bonding abilities. Cytosine, thymine and uracil may be modified at positions N1, C2, C4, C5, or C6 and still maintain their hydrogen bonding abilities.

10 Some base analogs have different hydrogen bonding attributes than the naturally occurring bases. For example, 2-amino-2'-dA forms three (3), instead of the usual two (2), hydrogen bonds to thymine (T).

15 Examples of base analogs that have been shown to increase duplex stability include, but are not limited to, 5-fluoro-2'-dU, 5-bromo-2'-dU, 5-methyl-2'-dC, 5-propynyl-2'-dC, 5-propynyl-2'-dU, 2-amino-2'-dA, 7-deazaguanosine, 7-deazadenosine, and N2-Imidazolylpropyl-2'-dG. For purposes of illustration, several base analogs are shown in Figures 5A and 5B.

20 Pendant Groups

A "pendant group" may be linked to the oligonucleotide. Pendant groups serve a variety of purposes which include, but are not limited to, increasing cellular uptake of the oligonucleotide, enhancing degradation of the target
25 nucleic acid, and increasing hybridization affinity. Pendant groups can be linked to any portion of the oligonucleotide but are commonly linked to the end(s) of the oligonucleotide chain. Examples of pendant groups include, but are not limited to: acridine derivatives (i.e.
30 2-methoxy-6-chloro-9-aminoacridine); cross-linkers such as psoralen derivatives, azidophenacyl, proflavin, and azidoproflavin; artificial endonucleases; metal complexes such as EDTA-Fe(II), o-phenanthroline-Cu(I), and porphyrin-Fe(II); alkylating moieties; nucleases such as amino-1-
35 hexanolstaphylococcal nuclease and alkaline phosphatase;

terminal transferases; abzymes; cholesteryl moieties;
lipophilic carriers; peptide conjugates; long chain
alcohols; phosphate esters; amino; mercapto groups;
radioactive markers; nonradioactive markers such as dyes;
and polylysine or other polyamines.

In one example, the oligonucleotide comprises an
oligonucleotide conjugated to a carbohydrate, sulfated
carbohydrate, or glycan. Conjugates may be regarded as a
way as to introduce a specificity into otherwise unspecific
DNA binding molecules by covalently linking them to a
selectively hybridizing oligonucleotide.

Cellular Uptake

To enhance cellular uptake, the oligonucleotide may be
administered in combination with a carrier or lipid. For
example, the oligonucleotide may be administered in
combination with a cationic lipid. Examples of cationic
lipids include, but are not limited to, lipofectin, dotma,
dope, and Dogs. The oligonucleotide may also be
administered in combination with a cationic amine such as
poly (L-lysine). Oligonucleotide uptake may also be
increased by conjugating the oligonucleotide to chemical
moieties such as transferrin and cholesteryls. In
addition, oligonucleotides may be targeted to certain
organelles by linking specific chemical groups to the
oligonucleotide. For example, linking the oligonucleotide
to a suitable array of mannose residues will target the
oligonucleotide to the liver.

The cellular uptake and localization of oligonucleotides
may be monitored by using labeled oligonucleotides.
Methods of labeling include, but are not limited to,
radioactive and fluorescent labeling. Fluorescently
labeled oligonucleotides may be monitored using
fluorescence microscopy and flow cytometry.

The efficient cellular uptake of oligonucleotides is well

established. For example, when a 20 base sequence phosphorothioate (PS) oligonucleotide was injected into the abdomens of mice, either intraperitoneally (IP) or intravenously (IV). The highest concentrations of oligonucleotide accumulated in the kidney and liver, with only very small amounts being found in the brain. Chain-extended oligonucleotides were also observed. Argrawal, S., et al. (1988) Proc. Natl. Acad. Sci. U.S.A. 85:7079-7083. When the PS 27-oligonucleotide α rev was given IV or to rats, the initial $T_{1/2}$ (transit out of the plasma) was 23 min, while the $T_{1/2\beta}$ of total body clearance was 33.9 hours. The long β half-life of elimination demonstrates that dosing could be infrequent and still maintain effective, therapeutic tissue concentrations. Iverson, P. (1991) Anti-Cancer Drug Des. 6:531.

The efficacy of oligonucleotide therapy is also well established. For example, when a 24-base sequence PS oligonucleotide targeted to human c-myc mRNA was infused, through a miniosmotic pump, into scid mice bearing the human K562 chronic myeloid leukemia cell line, mean survival times of the mice treated with the antisense oligonucleotides were six- to eightfold longer than those of mice untreated or treated with the sense controls or treated with an oligonucleotide complementary to the c-kit proto-oncogene mRNA. Furthermore, significantly less tumor burden in the brain and ovary was observed histologically compared with the controls. After injecting IP 3'-PS-modified chimeric oligonucleotides that were complementary to the initiation codon region of the NF- κ B mRNA (p65), a complete tumor involution was observed in 13 out of 13 antisense-treated mice. Untreated or sense-treated mice died by 12 weeks, whereas the treated animals had no recurrence for at least 5 months. Ratajczak, et al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89:11823.

This invention provides an antisense oligonucleotide or analog thereof comprising 10 or more contiguous bases or

base analogs from the sequence of bases of sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

1 This invention also provides an antisense oligonucleotide
or analog thereof comprising a sequence having 90% of
5 greater identity to sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

10 This invention also provides an antisense oligonucleotide or analog thereof comprising a sequence having 85% of greater identity to sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

This invention also provides an antisense oligonucleotide or analog thereof comprising a sequence having 80% of greater identity to sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

15 This invention also provides an antisense oligonucleotide or analog thereof comprising a sequence having 75% of greater identity to sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

20 This invention also provides an antisense oligonucleotide or analog thereof comprising a sequence having 70% of greater identity to sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

25 This invention further provides an antisense oligonucleotide or analog thereof comprising nucleotide sequence A, B, C, D, E, F, G, H, I, J, K, L, or M of Figure 1.

30 This invention also provides the above-described antisense oligonucleotides, wherein the nucleotide sequence comprises nucleotide sequence A, A', B, C, C', D, E, E', F, G, G', H, H', I, I', J, K, K', L, L', M, or M' of Figures 2A and 2B.

This invention further provides the above-described antisense oligonucleotides, wherein the oligonucleotide is conjugated to a peptide.

5 This invention also provides the above-described antisense oligonucleotides, wherein the oligonucleotide is encapsulated in a liposome or nanoparticle.

This invention also provides the above-described antisense oligonucleotides, wherein the phosphate backbone comprises phosphorothioate bonds.

10 In addition, this invention provides the above-described antisense oligonucleotides, wherein the backbone is bonded to one or more lipid substituents.

15 This invention also provides the above-described antisense oligonucleotides, wherein one or more of the oligonucleotides's sugars contain an -OMe group at their 2' positions.

This invention further provides the above-described antisense oligonucleotides, wherein the phosphate backbone consists essentially of phosphorothioate bonds.

20 This invention also provides the above-described antisense oligonucleotides, wherein the phosphorothioate is stereo regular.

25 This invention also provides the above-described antisense oligonucleotides, wherein the oligonucleotide is linked to an intercalating agent, a cross-linker, an endonuclease, a lipophilic carrier, an alkylating agent, a coordination complex, or a peptide conjugate, or a combination thereof

30 In addition, this invention provides the above-described antisense oligonucleotides, wherein the oligonucleotide is modified to reduce its ionic charge or increase its

hydrophobicity.

In an embodiment, an alkyl group is attached to increase the hydrophobicity of the oligonucleotide. In another embodiment, one or more oxygen atoms are replaced by sulphur atoms to decrease the hydrophobicity of the oligonucleotide.

This invention also provides the above-described antisense oligonucleotides, wherein the oligonucleotide comprises one or more short chain alkyl structures that replace some of the oligonucleotide's phosphodiester bonds.

This invention further provides the above-described antisense oligonucleotides, wherein the oligonucleotide is linked to one or more cholesteryl moieties.

This invention also provides the above-described antisense oligonucleotides, wherein the oligonucleotides comprises one or more bases with a C-5 propynyl pyrimidine modification.

The above-described antisense oligonucleotide wherein the oligonucleotide comprises one or more bases with a C-5 propynyl pyrimidine modification.

This invention provides a method of treating cancer, comprising introducing into a tumor cell an effective amount of the antisense oligonucleotide of the above-described antisense oligonucleotide wherein the oligonucleotide comprises one or more bases with a C-5 propynyl pyrimidine modification, thereby reducing the levels of bcl-2 protein produced and treating cancer.

In an embodiment, the cancer is epithelial cancer. In a further embodiment, the epithelial cancer is prostate cancer. In still a further embodiment, the epithelial cancer is lung cancer, in still a further embodiment, the

epithelial cancer is bladder cancer.

In a separate embodiment of the above method, the introducing comprises using a lipid as a delivery agent. In a further embodiment, the introducing comprises using porphyrin or lipofectin as a delivery agent.

In another embodiment, the effective amount is between 0.1 μM and 10 μM . In a further embodiment, the effective amount is between 0.1 μM and 4 μM . In a further embodiment, the effective amount is between 0.4 μM and 1 μM .

In addition, this invention provides a method of treating cancer, comprising introducing into a tumor cell an effective amount of the the above-described antisense oligonucleotide, thereby reducing the levels of bcl-xL protein produced and treating cancer. In an embodiment, the cancer being treated is epithelial cancer.

The actual effective amount will be based upon the size of the oligonucleotide, the biodegradability of the oligonucleotide, the bioactivity of the oligonucleotide and the bioavailability of the oligonucleotide. If the oligonucleotide does not degrade quickly, is bioavailable and highly active, a smaller amount will be required to be effective. The effective amount will be known to one of skill in the art; it will also be dependent upon the form of the oligonucleotide, the length of the oligonucleotide and the bioactivity of the polypeptide. One of skill in the art could routinely perform empirical activity tests to determine the bioactivity in bioassays and thus determine the effective amount.

This invention also provides the above-described methods, wherein the effective amount is between 0.1 μM and 10 μM . In an embodiment of the above-described method, the effective amount is between 0.1 μM and 4 μM .

In addition, this invention provides the above-described methods, wherein the effective amount is between 0.4 μM and 1 μM .

5 This invention further provides the above-described methods, wherein the cancer is epithelial cancer.

This invention also provides the above-described methods, wherein the epithelial cancer is prostate cancer.

This invention also provides the above-described methods, wherein the epithelial cancer is lung cancer.

10 In addition, this invention provides the above-described methods, wherein the epithelial cancer is bladder cancer.

This invention also provides the above-described methods, wherein the introducing comprises using a lipid as a delivery agent.

15 This invention further provides the above-described methods, wherein the introducing comprises using porphyrin or lipofectin as a delivery agent.

20 This invention also provides the above-described methods, wherein the effective amount is between 0.1 μM and 10 μM .
In an embodiment of the above-described method, the effective amount is between 0.1 μM and 4 μM .

In addition, this invention provides the above-described methods, wherein the effective amount is between 0.4 μM and 1 μM .

25 This invention also provides a method of promoting the regression of vascular lesions, comprising introducing into a vascular cell an amount of the the above-described antisense oligonucleotides effective to reduce the levels of bcl-xL protein produced, thereby promoting the

regression of vascular lesions.

This invention further provides the above-described methods, wherein the introducing comprises using a lipid as a delivery agent.

5 This invention also provides the above-described methods, wherein the introducing comprises using porphyrin or lipofectin as a delivery agent.

10 This invention also provides the above-described methods, wherein the effective amount is between 0.1 μ M and 10 μ M. In an embodiment of the above-described method, the effective amount is between 0.1 μ M and 4 μ M.

In addition, this invention provides the above-described methods, wherein the effective amount is between 0.4 μ M and 1 μ M.

15 This invention also provides a pharmaceutical composition comprising an effective amount of any of the above-described antisense oligonucleotides or analogs thereof and a pharmaceutically acceptable carrier.

20 Pharmaceutically acceptable carriers are well known to those skilled in the art and include, but are not limited to phosphate buffer and saline. Additionally, such pharmaceutically acceptable carriers may be aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, 25 Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers such as 30

those based on Ringer's dextrose, and the like. Preservatives and other additives may also be present, such as, for example, antimicrobials, antioxidants, chelating agents, inert gases and the like.

5 This invention further provides the above-described pharmaceutical compositions, wherein the effective amount is between 0.1 μ M and 4 μ M. In an embodiment of the above-described composition, the effective amount is between 0.1 μ M and 4 μ M.

10 In addition, this invention provides the above-described pharmaceutical compositions, wherein the effective amount is between 0.4 μ M and 1 μ M.

15 This invention also provides the above-described pharmaceutical compositions, wherein the oligonucleotide is encapsulated in a liposome or nanoparticle.

This invention further provides the above-described pharmaceutical compositions, wherein the pharmaceutical composition comprises tetra meso-(4-methylpyridyl)porphine or tetra meso-(anilinium)porphine or a combination thereof.

20 This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which
25 follow thereafter.

EXPERIMENTAL DETAILS

First Series of Experiments

30 Forty 18- and 20-mer mixed phosphate/phosphorothioate backbone oligonucleotides with or without C-5-propyne pyrimidine modifications chosen by "walking" along the

bcl-xL mRNA have been evaluated as to their ability to decrease bcl-xL protein expression in human prostate cancer cell lines (LNCaP, DU145 and PC3). The cell lines were obtained from the ATTC (Rockville, MD). The cells were grown in RPMI 1640 media supplemented with 10% FBS in 5% CO₂ atmosphere and routinely passaged when 90-95% confluent. All oligonucleotides were at least 95% full-length material when analysed by a reverse-phase HPLC.

Two novel agents for the intracellular delivery of oligonucleotides were used: tetra meso-(4-methylpyridyl)porphine (TMP) and tetra meso-(trimethylammonium)porphine (TAP). TMP and TAP both are cations which form 2:1 and 3:1 complexes with oligonucleotides. Human prostate carcinoma cells were plated in 6-well tissue culture dishes and next day (when 80-90% confluent) were washed once with OPTI-MEM and treated with prepared complex of oligonucleotide (1 μ M) plus TMP (2 μ M) or TAP (3 μ M) for 4 h at 37°C. The cells were then washed once with RPMI-10% FBS and were allowed to recover in RPMI-10% FBS for another 20 h. At this time cells were washed twice with PBS and then protein and/or mRNA were extracted and analysed.

Western blot analysis demonstrated the effective down-regulation of bcl-xL protein with five of the most active oligonucleotides (T31057, T31061, T31062, T31065, T31067). For the western blot analyses approximately 10⁶ cells were lysed for 30 min on ice in 100 μ l of RIPA buffer (50 mM Tris-HCl, pH 8.0 - 150 mM NaCl - 0.1% SDS - 1% NP40 - 0.5% sodium deoxycholate). Aliquots of cell extracts containing 20 μ g of protein were electrophoresed in 12% SDS-polyacrylamide gel and transferred to PVDF membrane (Millipore Corp., Bedford, MA). Filters were blocked and stained with an 1:200 dilution of rabbit anti-bcl-x polyclonal antibody (Santa Cruz Biotechnology Inc., Santa-Cruz, CA). After washing, filters were incubated in a 1:10,000 dilution of peroxidase-conjugated anti-rabbit

secondary antibody (Amersham Life Sciences, Arlington Heights, IL). ECL was performed using the procedure recommended by the manufacturer and the filters were exposed to X-ray film.

5 Control oligonucleotides did not affect expression of bcl-xL and cell viability. The most active oligonucleotides are complementary to regions on the 3'-end of the bcl-xL mRNA. Moreover, C-5 propyne pyrimidine modified oligonucleotides seem to be more effective than
10 pyrimidine unmodified.

These five active oligonucleotides (T31057, T31061, T31062, T31065, T31067) alone did not induce apoptosis in DU145 prostate cancer cell lines, as is shown with DAPI-staining. To determine the amount of apoptotic nuclei, cells were
15 plated in 4-chamber tissue culture slides (Nunc, Inc., Naperville, IL) and after the experiment, were washed with PBS, fixed with 90% ethanol/ 5% acetic acid, and after 2 rinses with PBS, stained with a 1.5 mg/ml solution of DAPI in PBS. The slides were washed twice with PBS, mounted and
20 photographed using a Nikon phase-fluorescence microscope.

In an attempt to approximate clinical drug resistance in prostate cancer a commonly used prostate cancer cell line, LNCaP, was genetically manipulated so that these cells would overexpress the human bcl-xL protein. This cell line
25 is a popular model for the study of prostate cancer, because it retains some of the most prominent differentiated features of the human prostate cell. LNCaP cells also have proven to be growth responsive to androgen steroids in vitro. This bcl-xL overexpressing cell line
30 provides a useful in vitro model for the subsequent study of drug resistant adenocarcinoma of the prostate gland.

The human prostate cancer cell line LNCaP was received from ATTC (Rockville, MD). LNCaP cells were propagated as monolayer culture in RPMI 1640 supplemented with 10% FBS,

in 5% CO₂ atmosphere. Confluent cultures were routinely passaged approximately weekly, with RPMI 1640, containing 5 mM EDTA.

5 LNCaP cells transfected with the neomycin-selectable pSFFV/bcl-xL plasmid or with a control, neomycin-resistant expression vector pSFFV. Aliquots containing 10 µg of plasmid and 5 µg of lipofectin reagent (Life technologies, Inc., Gaithersburg, MD) in serum-free OPTI-MEM (Life Technologies, Inc.) were added to cultured cells. The
10 transfection media was replaced 4 h later with RPMI-10% FBS medium. Individual colonies were selected from these plates after approximately 3-4 weeks of routine maintenance in RPMI-10% FBS containing 0.6 mg/ml G418 sulfate (Geneticin, Life Technologies, Inc.). The bcl-xL-transformed LNCaP
15 cells were cultured in RPMI-10% FBS, supplemented with 0.3 mg/ml G418 sulfate in 5% CO₂ atmosphere.

Two clones of LNCaP cells overexpressing bcl-xL protein (1072-4 and 1072-5) have been obtained after transfection of wild type LNCaP cells with the plasmid vector
20 pSFFV/bcl-xL and lipofectin. Also a mock transfectant clone of LNCaP cells carrying neo[®] resistance gene (1072-3) was used for the control experiments. Clone 1072-4 demonstrates 10-fold overexpression, and clone 1072-5 - 4-fold overexpression of bcl-xL protein. Western blot analysis for
25 bcl-xL protein was performed as described above. Results for bcl-xL protein expression were confirmed by Northern blot analysis for bcl-xL mRNA expression, demonstrating significant elevation of this mRNA in bcl-xL transformed cell lines. For the Northern blot analysis, the total RNA
30 was isolated from the cells using TRIZOL reagent (GIBCO BRL), and 20 µg aliquotes were separated in RNA-formaldehyde gel, blotted onto nylon membranes (Schleicher & Schull), UV-linked and prehybridized for two hours at 42 °C in the standard hybridization solution. Then
35 the blot was hybridized overnight with the PCR-amplified fragment of human bcl-xL cDNA at 42 °C. Bcl-xL coding

fragment was amplified from pSFFV/bcl-xL plasmid using bcl-x specific primers. The primer sequences were: bcl-x-upstream, 5'-ATGTCTCAGAGCAACCGGGA-3'; and bcl-x-downstream, 5'-TCATTTCCGACTGAAGAGTG-3'. Twenty five cycles of amplification were performed in DNA Thermal Cycler (Perkin-Elmer, Norwalk, CT) at 94 °C (30 sec), 55 °C (30 sec), and 72 °C (30 sec). The PCR products were analysed on a 1.2% agarose gel. The resultant fragment was labeled by random primer method to the specific activity 10⁷ cpm/ng of the probe and used for the hybridization. After washings blots were autoradiographed for 24h at -80 °C. Blots were stripped of radioactivity and reprobbed with a ³²P-labeled G3PDH probe to confirm the equal loading.

Overexpression of bcl-xL protein does not affect the relative growth rate of LNCaP cells in culture medium containing 10% FBS as determined by MTT assay. For this assay LNCaP cells were seeded in 24-well tissue culture plates. Next day after the experiment the old media was removed, the cells were washed once with PBS, and RPMI containing 0.5 mg/ml MTT (Sigma Co, St. Louis, MO) was added. The cells were incubated at 37°C for 4 h, then solubilization solution (0.04N HCl in isopropyl alcohol) was added and optical density was measured at 540 nm.

Clonal derivatives of bcl-xL transfected LNCaP cells that were overexpressing high amounts of bcl-xL protein were resistant to taxol (up to 100 nM) and mitoxantrone (up to 4 µM) treatment for 24 h. These drugs have been shown to induce apoptosis in parental LNCaP cells or neomycin-expressing control-transfected LNCaP cells. This finding strongly supports the hypothesis that bcl-xL gene product is a factor in the development of chemotherapy resistance in hormone-refractory prostate cancer.

Antisense oligonucleotides provide an efficient means for bcl-xL protein elimination in prostate cancer cell lines. When antisense oligonucleotides were applied to the bcl-xL

overexpressor, the down-regulation of this anti-apoptotic protein was not apparant. A possible way to down-regulate bcl-xL protein in such case could be using antisense oligonucleotide therapy in combination with
5 chemotherapeutic drugs.

It has been shown recently (Am J Pathol 1996, 148:1567-1576) that expression of several anti-apoptotic members of the bcl-2 family, including bcl-2 and bcl-xL proteins increases during progression of prostate cancers.
10 Our transfected clones overexpressing bcl-xL can mimic the clinical model of adenocarcinoma of the prostate gland. Demonstration of the mechanism of action of different drugs and their combinations with antisense oligonucleotides in these transfected LNCaP cells could help form the basis for
15 a better understanding of prostate cancer.

Forty modified oligonucleotides complementary to the bcl-xL mRNA have been evaluated as to their ability to decrease bcl-xL protein expression in human prostate cancer cell lines (LNCaP, DU145, and PC3).

20 Two novel agents for the intracellular delivery of oligonucleotides were used: tetra meso(4-methylpyridyl)porphine (TMP) and tetra meso(trimethylammonium)porphine (TAP).

Western blot analysis demonstrated effective
25 down-regulation of bcl-xL protein with five of the most active oligonucleotides (T31057, T31061, T31062, T31065, T31067).

These active oligonucleotides alone did not induce apoptosis in DU145 cells.

30 Two clones of LNCaP cells overexpressing bcl-xL protein and clone carrying neo resistance gene only have been obtained after transfection of wild type LNCaP cells with

the plasmid vectors pSFFV/bcl-xL and pSFFV.

Overexpression of bcl-xL protein and mRNA has been shown by western and northern blot analyses.

Overexpression of bcl-xL protein does not affect the relative growth rate of LNCaP cells in culture medium containing 10% FBS.

Bcl-xL overexpressing clones of LNCaP cell lines demonstrated high resistance to taxol (100 nM) and mitoxantrone (4 μ M) treatment.

Active antisense oligonucleotides do not down-regulate bcl-xL protein in the LNCaP clones overexpressing bcl-xL.

Second Series of Experiments

Bcl-xL in Prostate and Bladder Cancer Cells: Chemosensitization After Downregulation by Treatment with Directed ("Antisense") Oligonucleotides

Introduction

This study demonstrated the role of bcl-xL and bcl-2 proteins for the regulation of the sensitivity of cancer cells to different drugs, including taxanes and antracyclines. In conclusion, it was demonstrated that treatment with bcl-xL antisense oligonucleotides induces sensitization of prostate and bladder carcinoma cell lines to chemotherapy treatment, suggesting that Bcl-xL provides one of the defending mechanisms for cancer cells allowing them to tolerate chemotherapeutic treatment.

Several pathways have been described that regulate programmed cell death (apoptosis). One these implicates bcl-2 family proteins as critical in apoptosis commitment. The bcl-2 family includes several homologous proteins that

may be either pro- or antiapoptotic. Bcl-2 and bcl-xL are antiapoptotic, whereas bax promotes apoptosis. Together, these proteins form a complex network of heterodimers and homodimers that govern the relative sensitivity of a cell to potentially apoptotic stimuli.

Bcl-x is a relatively new member of the bcl-2 family of apoptosis-related proteins, and its expression has been detected in a range of normal tissues, particularly in the central nervous system and thymus. Immunocytochemical studies have demonstrated that the bcl-x protein can be detected in numerous tumor cell lines as well. Expression has also been found in the epithelial cells of the normal prostate (Krajewski, et al., Cancer Res 1994; 54: 5501-5507), where, it is speculated, it contributes to the hormone-dependent control of programmed cell death. Krajewski, et al., (1996) found that 64/64 (100%) of cases of adenocarcinoma of the prostate stained positively for bcl-x protein. Staining intensity seemed to be correlated with increasing Gleason score and the presence of metastases. Bcl-xL protein expression has also been demonstrated in 50% of clinical specimens of low stage bladder tumors (Gazzaniga, Oncol. Rep. 1998 5,901-04)

Since bcl-xS appears not to be expressed either in prostatic tumors, or in prostate cell lines in tissue culture, this staining most likely originates from bcl-xL. This 25.6 kDa protein has been shown in a number of cell lines to be a potent protector of cellular apoptosis induced by antineoplastic agents (Kim, 1997). When transferred into the murine IL-3-dependent prolymphocytic line FL5.12, bcl-xL greatly reduced the pro-apoptotic effects of bleomycin, cisplatin, etoposide, vincristine and doxorubicin (Ibrado, 1997; Minn, et al., 1995). In U937 cells, inhibitory effects on camptothecin-induced apoptosis were shown to be dependent on the intracellular concentration of bcl-xL protein. Diminution of cell death in response to etoposide, vinblastine, paclitaxel and

5 cisplatinum were also observed (Schmitt, Exp. Cell Res. 1998 240 107-21). Bcl-xL can also block cellular apoptosis in settings in which bcl-2 is ineffectual (Gottschalk, PNAS 1994 91 7350). Several agents, such as butyrate in human
10 fibroblasts (Chung, Rad Res. 149, 187 (1998), and paclitaxel in LNCaP and PC3 human prostate cancer cells (Liu) induce apoptosis that can be correlated with substantial decreases in the expression of bcl-xL protein. In the latter case, this occurred in the absence of any decrease in bax, bak, fas or fas ligand expression.

15 Bcl-2 protein expression has been found in 66% of low stage bladder tumors and 100% of high stage tumors. Expression is also found quite extensively in primary prostate cancer specimens (McConnell, 1992), although it has been observed in only 33% of samples of prostate tumors obtained from the bone marrow of patients with hormone-refractory disease (McDonnell J. Urol. 1997 569-574). At the present time, it appears that bcl-xL and bcl-2 have similar antiapoptotic function, and it is not clear why both are expressed in at
20 least some prostate cancer cells. Notwithstanding, based on these data it is possible that overexpression of bcl-xL and bcl-2 protein might be factors enabling prostate cancer cells to survive in an androgen-deprived environment.

Materials and Methods

25 Reagents

Taxol and etoposide are products of Bristol-Myers Squibb, Princeton, NJ. Taxotere is a product of Rhone-Poulenc Rorer, Collegeville, PA. Polysorbitan-80 and EL-cremophor, the carriers to taxotere and taxol, respectively, were
30 obtained from Sigma (Milwaukee, WI). Mitoxantrone was from Immunex (Seattle, WA), and vinblastine was from Eli Lilly (Indianapolis, IN)

Cell Culture and Transformation of Cell Lines with Bcl-xL Plasmid

Human PC3 and LNCaP prostate cancer cell lines, and the T24 bladder carcinoma cell line were obtained from American Type Culture Collection (Rockville, MD). They were grown in RPMI 1640 medium (Gibco BRL, Grand Island, NY), containing 10% (v/v) heat inactivated (56°C) fetal bovine serum (FBS) (Gibco BRL, Grand Island, NY), supplemented by 1% non-essential amino acids, 1% pyruvate, and 100 units/mL penicillin G sodium and 100 mg/mL streptomycin sulfate. All cell lines were cultured in 5% CO₂ atmosphere and were routinely passaged when 90-95% confluent.

LNCaP cells were transfected with the neomycin-selectable pSFFV/bcl-xL plasmid (obtained from Dr. Stanley Korsmeyer, Washington University School of Medicine, St. Louis, MO) or with a control, neomycin-resistant expression vector pSFFV. Aliquots containing 10 µg of plasmid and 5 µg of Lipofectin, reagent (Life technologies, Inc., Gaithersburg, MD) in serum-free OPTI-MEM (Life Technologies, Inc.) were added to a cultured cells. The transfection media was replaced 4 h later with RPMI-10% FBS medium. Individual colonies were selected from these plates after approximately 3-4 weeks of routine maintenance in RPMI-10% FBS containing 0.6 mg/ml G418 sulfate (Geneticin, Life Technologies, Inc.). The bcl-xL-transformed LNCaP cells were cultured in RPMI-10% FBS, supplemented with 0.3 mg/ml G418 sulfate in 5% CO₂ atmosphere.

Oligonucleotides

Mixed-phosphate backbone (phosphorothioate and phosphodiester) oligonucleotides with or without C-5 propynyl modified pyrimidines, complementary to different regions of bcl-xL mRNA were obtained from The Aronex Pharmaceuticals Inc. (Woodland, Texas). All oligonucleotides were 95% full-length material when analysed by a reverse-phase HPLC.

Treatment of cells with oligonucleotide-Lipofectin, or oligonucleotide-porphyrin complexes

Prostate and bladder carcinoma cells were seeded the day before experiment in 6-well plates at density 3×10^5 cells per well (to be 80-90% confluent on the day of the experiment).

Oligonucleotides were delivered to the cells in the form of complexes with cationic lipid Lipofectin, (GIBCO BRL, Gaithersburg, MD) or cationic porphyrins. In the first case equal volumes of oligonucleotide (5 mM) and Lipofectin, (5 mg) in reduced serum OPTI-MEM medium (GIBCO BRL, Gaithersburg, MD) were mixed and allowed to form a complex for 30 min at room temperature. The mixture was diluted 1:9 with OPTI-MEM media and added to T24 cells rinsed with OPTI-MEM medium. The final concentrations were 1mM for oligonucleotide and 5 mg/ml for Lipofectin. Control cells were treated with equivalent concentration of Lipofectin, or medium alone. Cells were incubated at 37°C for 5 hours, then medium was replaced for McCoy 5A supplemented with 10% FBS.

For cationic porphyrin delivery, oligonucleotide was premixed with TAP or TMP in OPTI-MEM medium to the final concentrations described in the Results for each cell line. Complex was formed for 15 min at room temperature, cells were washed once with OPTI-MEM and treated with prepared complex for 4 h at 37°C. Then cells were washed once with RPMI-10% FBS and were allowed to recover in RPMI-10% FBS for another 20 h. At this time cells were washed twice with PBS and then protein and/or mRNA were extracted as described and analyzed.

Western Blot Analysis

Cells treated with oligonucleotide - porphyrine or oligonucleotide - Lipofectamin, complex were washed twice with cold PBS and lysed on ice for 30 min in 100 ml of cold

RIPA buffer [50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 0.1% SDS, 1% NP40, and 0.5 % sodium deoxycholate] with freshly added 0.1 mg/ml phenylmethylsulfonyl fluoride, 1 mM sodium orthovanadate, and 1 mg/ml aprotinin. Cell debris were removed by centrifugation at 14,000g for 10 min at 4°C. Protein concentrations were determined using the Bio-Rad protein assay system (Bio-Rad Laboratories, Richmond, CA).

Aliquots of cell extracts containing 20-50 mg of total protein were resolved in 12% SDS-PAAG and transferred to Immobilon-P PVDF membranes (Millipore Corp., Bedford, MA). Filters were blocked for 1 h at room temperature in Blotto A [5% nonfat milk powder in TBS-T: 10 mM Tris-HCl (pH 8.0), 150 mM NaCl, 0.05% Tween 20], and then incubated for 1 h at room temperature in Blotto A containing a 1:200 dilution of either rabbit anti-Bcl-xL, anti-Bax, or anti-Bak polyclonal antibodies or mouse anti-Bcl-2 mAb (all from Santa Cruz Biotechnology Inc., Santa Cruz, CA). After washing in TBS-T buffer (3x5 min, room temperature), filters were incubated for 45 min at room temperature in Blotto A containing a 1:10,000 dilution of corresponding peroxidase conjugated anti-rabbit or anti-mouse secondary antibody (Amersham, Arlington Heights, IL). After washing in TBS-T, ECL was performed according the recommendation of the manufacturer.

RNA Isolation and Northern Blot Analysis.

Total RNA was isolated from 5×10^7 cells using RNeasy Mini kit from QIAGEN (Santa-Clarita, CA). Forty μ g of total RNA were electrophoresed on a 1% denaturing agarose gel containing formaldehyde, transferred to a Nytran-Plus nylon membranes (Schleicher&Schuell, Keene, NH) and covalently bound to a membrane by UV-irradiation. A bcl-xL coding region cDNA fragment generated by PCR and 32 P - labeled by random primer method was hybridized to the filter-bound RNA in 50% formamide, 5xSSC, 5xDenhardt solution, 0.5% SDS and 0.1 mg/ml sonicated salmon sperm DNA overnight at 42°C. The

membrane was washed 3 times for 15 min in 1xSSC, 0.1% SDS at 42oC, and 3 times for 10 min in 0.1xSSC, 0.1% SDS at 65oC. The filter was exposed to Kodak X-ray film with intensifying screens for 3 days at -70oC and developed.

5 MTT assay for determination of cell viability.

The cytotoxicity of the combinations of drugs with oligonucleotides was determined by measurement of cell viability by use of colorimetric MTT assay.

10 LNCaP cells were seeded in 48-well tissue culture plates and treated next day subsequently with oligonucleotides and drugs at indicated concentration as described in Results. Then drug-contained media was removed, and the cells were washed once with PBS. Corresponding culture media (RPMI for prostate and McCoy 5A for bladder cancer cells) containing 0.5 mg/ml MTT (Sigma Co, St. Louis, MO) was added to each well. The cells were incubated at 37oC for 4 h, then equal amount of solubilization solution (0.04N HCl in iso-propyl alcohol) was added to each well and mixed thoroughly to dissolve the crystals of MTT formazan. After 15 all crystals were dissolved, the plates were read on a Dynatech MR600 Microplate Reader, using a wavelength of 540 nm. 20

Results

25 PS-PO Oligonucleotides directed against Bcl-xL mRNA.

In the present study we have used an antisense strategy for the down-regulation of bcl-xL in prostate cancer cell lines. Forty oligonucleotides (twenty 20-mers and twenty 18-mers) have been chosen by the "walking along the mRNA" method. An arrangement of the oligonucleotides on Bcl-xL mRNA is shown on Scheme 1, where numbers reflected the base positions of the oligonucleotides. Each oligonucleotide has been assigned a number for further convenience and will be 30

mention in the text according to this number. Oligonucleotides have mixed phosphorothioate-phosphodiester backbone. Oligonucleotides with even numbers have regular heterocyclic bases whereas odd numbers
5 oligonucleotides included C-propynyl modified cytosin and tymidine bases. Oligonucleotides were screened for their antisense activity to down-regulate Bcl-xL protein expression using Western blot analysis.

Optimal conditions for the delivery of oligonucleotides
10 within the cells

To deliver oligonucleotides into cells we have used two different types of the delivery agents: lipids, such as Lipofectin,, and cationic porphyrins (tetra meso-
15 (4-methylpyridyl)porphine (TMP) and tetra meso-(trimethylammonium)porphine (TAP)).

Lipofectin, has been used to transfect T24 cells with antisense oligonucleotides. The optimal condition for the delivery was 5 mg/ml Lipofectin, and 0.5 μ M of oligonucleotide in reduced serum OPTI-MEM medium.
20 Unfortunately, with prostate cancer cell lines Lipofectin, did not demonstrate out good results because of high cytotoxicity. To deliver oligonucleotides into LNCaP and PC-3 cell line cationic porphyrins have been used. Porphyrins are novel class of agents for the intracellular
25 delivery of oligonucleotides. They have been investigated by confocal microscopy and flow cytometry methods. It has been demonstrated that they form complexes with oligonucleotides and deliver them to the cell nucleus, where the porphyrin had completely dissociated from the
30 oligonucleotide. No non-specific cytotoxic effects has been observed after the using of the porphyrins as delivery agents. Porphyrins demonstrated different activity in different cell lines. The optimal conditions for LNCaP cells were 3 μ M of TAP and 1 μ M of oligonucleotide; for

PC-3 cells - 7 μ M of TMP and 2 μ M of oligonucleotide; and for T24 cells - 9 μ M of TMP and 3 μ M of oligonucleotide. After cell treatment with the oligonucleotides, total protein was harvested and level of Bcl-xL protein expression was determined comparing to the untreated cells.

Evaluation of Bcl-xL protein levels by Western blotting

The Bcl-xL protein was detected as a doublet with a molecular weight ranging from 29-31 kDa, a finding consistent with previous reports. During the screening of forty oligonucleotides, treatment with several sequences produced significant reductions in the steady-state levels of Bcl-xL protein expression. Compare to the untreated cells or cells treated with inactive oligonucleotides, the level of Bcl-xL expression decreases to 5-30% in case of the most active oligonucleotides. Some sequences provide moderate reduction of the Bcl-xL expression. A 19 kDa band which would correspond to the Bcl-xS protein was not detected in studied cells. Level of actin (control protein) remains unchanged all the times (Data are not shown). The oligonucleotide treatment also did not cause reductions in the total amount of protein, nor did it inhibit the growth of these cells, as was demonstrated by MTT assay.

Down-Regulation of Bcl-xL Protein and mRNA Expression with PS-PO Oligonucleotides in Different Cell Lines

Screening of oligonucleotide set revealed several the most active oligonucleotides (numbers 29, 41, 43, 57, 61, 62 and 63). They decreased bcl-xL protein expression to 5-30% in tissue culture comparing to the untreated cells or cells, treated with oligonucleotides along, without delivery agent. To address the question of the mechanism of action of those oligonucleotides, we studied bcl-xL mRNA expression in treated and untreated cells by the Northern hybridization method, using bcl-xL cDNA fragment as a

probe. A putative antisense mechanism of action for the majority of the active oligonucleotides (29, 41, 43, 57 and 61) is supported by the fact of down-regulation of Bcl-xL mRNA level. Oligonucleotides 62 and 63 possess a G-quartet and are likely not specific. In our further experiments we were not using those non-specific oligonucleotides. Considering the position of the active oligonucleotides on the bcl-xL mRNA, it can be assumed that those oligonucleotides are complementary to the most open (and so the most accessible for the RNase H action) mRNA regions.

C-propyne-modified oligonucleotides potentiate bcl-xL down-regulation. The ability of oligonucleotides to down-regulate bcl-xL protein expression differs in different cell lines (Data are not shown). The efficiency of the oligonucleotide is highly dependent on the delivery agent. Identical oligonucleotides demonstrated different bcl-xL down-regulation when delivered with different agents. However, it has to be emphasized that if oligonucleotide is active, it is active in any cell line, with any delivery agent.

Since the ability of cell to undergo apoptosis is determined by the ratio between anti- and proapoptotic members of bcl-2 family, we investigated if the treatment of the cells with antisense bcl-xL oligonucleotides affects the other related proteins. Our results demonstrated that there is not a unique action of bcl-xL antisense oligonucleotides on bax and bcl-2 proteins. The result is highly dependent on the cell line, the delivery agent and on the oligonucleotide sequence (Table 1). Cationic porphyrin-mediated delivery appears to provide increased specificity of oligonucleotide action. Oligonucleotides 29, 41 and 43, when delivered with TMP caused down-regulation of both anti-apoptotic proteins bcl-xL and bcl-2, but upregulated the proapoptotic protein bax in all studied cell lines (however, oligo 29 does not show significant activity in the T24 bladder cell carcinoma

cell line with TMP delivery). It is suggested that treatment of the cells with oligonucleotides 29 and especially 41 and 43, can stimulate apoptotic processes within the cells and hence make cells more sensitive to chemotherapy treatment.

5 MTT assay demonstrated that treatment of the cells with oligonucleotides alone caused decrease of cell viability. Preliminary experiments with oligo 43 demonstrated a two-fold sensitization of the PC3 and T24 cell lines to treatment by mitoxantrone compare to the control (treated with drug only) cells. This effect is highly visible at low
10 drug concentration (less than 0.5 μ M) and less obvious at high concentrations of mitoxantrone. Although oligonucleotide 57 makes cells more sensitive to chemotherapy, the effect is less pronounced. Oligonucleotide
15 45 does not cause bcl-xL down-regulation and does not sensitize cells to mitoxantrone treatment.

Table 1

Regulation of Bcl-Family Proteins Expression with Chimeric PS-PO Oligonucleotides in Prostate and Bladder Cancer Cell Lines

	PC-3 Cell Line			LNCaP Cell Line			T24 Cell Line					
	TMP			TAP			lipofectin			TMP		
	bcl-xL	bax	bcl-2	bcl-xL	bax	bcl-2	bcl-xL	bax	bcl-2	bcl-xL	bax	bcl-2
29	↑	↑	↑	↑	↑	↑	↑	↑	↑	-	-	-
41	↑	↑	↑	↑	-	↑	-	↑	↑	↑	↑	↑
43	↑	↑	↑	↑	↑	↑	-	↑	-	↑	↑	↑
57	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	-
61	↑	↑	↑	↑	↑	↑	ND	ND	ND	ND	ND	ND
62	↑	↑	-	↑	↑	-	↑	↑	-	↑	↑	↑
62s	↑	↑	-	↑	↑	↑	-	↑	↑	↑	↑	-
63	↑	-	-	↑	-	-	-	-	-	↑	↑	↑

Marked reductions in Bcl-xL protein were detectable within 24 hours of additions of active oligonucleotides to LNCaP (60% reduction) and T24 cells (90% reduction), based on estimates using Western Blot analysis. Reductions of more than 80% in the relative levels of Bcl-2 protein in T24 cells treated with antisense oligonucleotides occurred also within 24 hours after treatment with antisense oligonucleotides. Thus, reductions in both proteins levels after antisense oligonucleotides treatment occur simultaneously. That allowed us to use an effect of oligonucleotide-mediated down-regulation of anti-apoptotic proteins Bcl-xL and Bcl-2 to sensitize cells to chemotherapy.

5

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000